

ਸ਼ਹੀਦ ਮੇਜਰ ਹਰਮਿੰਦਰਪਾਲ ਸਿੰਘ (ਸ਼ੈ.ਚੌ.) ਸਰਕਾਰੀ ਕਾਲਜ, ਸਾਹਿਬਜ਼ਾਦਾ ਅਜੀਤ ਸਿੰਘ ਨਗਰ

SHAHEED MAJOR HARMINDERPAL SINGH (Shaurya Chakra) GOVERNMENT COLLEGE, SAHIBZADA AJIT SINGH NAGAR

ਫੋਮ-6, ਸਾਹਿਬਜ਼ਾਦਾ ਅਜੀਤ ਸਿੰਘ ਨਗਰ (ਮੋਹਾਲੀ)-160056 Phone No. 0172-2225164 e-mail ID: principal_gcimohali@gmail.com



Research Papers Published by the Faculty in various National and International Journals

TABLE OF CONTENT: Research Papers Published By Faculty Members During Last Five Years

S.No	Title of the Paper	Author	Page no
1	Men as Oppressors and saviours, Women as Victims: An Analysis of Gender in Punjab 1984	Gurpreet Kaur	3-4
2	Sub classes of analytic function related to sigmoid function	Dr. Harjinder Singh	5
3	A new sub class of univalent functions	Dr. Harjinder Singh	6
4	Data mining : A Library utility model	Pardeep Rattan	7
5	Economic Dilemma and Suicide Among Farmers in Punjab	Vipan Chaudhary	8
6	Relationship between Social Support and Resilience among Farmers	Vipan Chaudhary	9
7	Investigating TiO ₂ at two different calcination temperatures	Shilpy Bhullar	10
8	A second order smoothing penalty Function Algorithm for constrained optimization problems	Dr. Amrit Pal Singh	11
9	Collaborative governance for urban sustainability: implementing solar cities	Rohit Barach	12
10	Rapid Green - Synthesis of TiO ₂ Nanoparticles for therapeutic applications	Shilpy Bhullar	13
11	Fekete-Szego Inequality for Certain Classes of Close to Convex Functions	Dr. Harjinder Singh	14
12	A smoothing technique for square root exact penalty function in constrained optimization	Dr. Amrit Pal Singh	15
13	Factors Influencing synthesis of titania nanoparticles- A Short review	Shilpy Bhullar	16
14	Titanium dioxide nanoparticles synthesized using different reagents	Shilpy Bhullar	17
15	- In-vitro pH-responsive released of imatinib from iron-supplement coated anatase TiO ₂ nanoparticles	Shilpy Bhullar	18
16	Coefficient Inequality for A Combined Subclass of Various Classes of Regular Functions	Dr. Harjinder Singh	19

2 | Page




Coordinator
NAAC
SMHS Government College
Sahibzada Ajit Singh Nagar



Principal
SMHS Govt. College
Sahibzada Ajit Singh Nagar

Men as Oppressors and saviours, Women as Victims: An Analysis of Gender in Punjab 1984 by Gurpreet Kaur



Coordinator
NAAC
SMHS Government College
Sahibzada Ajit Singh Nagar

Harjeet Goyal

Principal
SMHS Govt. College
Sahibzada Ajit Singh Nagar

1:09

Vol. 46
L1E2 47

Vol. No. 04, Issue No. 16, December- 2017

A UGC Approved Journal No. 63712

ISSN - 2395-1133 (Print)

Men as Oppressors and Saviours, Women as Victims: An Analysis of Gender in Punjab 1984

Author: Gurpreet Kaur, Assistant Professor in English
Government College, S A S Nagar, Mohali. Email: gurpreetgcmohali@gmail.com

The present research attempts to comprehend the portrayal of women in the film Punjab 1984. It investigates the women in the narrative for their opinion and voice. It underscores the stereotypical depiction of characters and the consequent victimhood of women. It questions the authority of men who possess the ability to oppress as well as rescue women.

Gender refers to the social system of differentiating between people on the basis of their sex. It recognises a collection of qualities and behavioural traits that the society expects from females or males. The idea of gender was delineated in order to stress the social formulation of masculinity and femininity, and social arrangement of relations between women and men. The distinction between sex and gender is one of the main concerns of feminist theory. In sex-gender dichotomy, sex refers to the biological distinction between men and women, and is viewed as an innate quality which distinguishes each one of us into the categories of male and female. Thus, gender implies the socially constructed characters, traits, and conduct that the society associates with men and women separately. Further, it can also be inferred that gender is not the direct result of sex, and not even as fixed as sex. An individual is born with a particular sex, and adopts the constructed norms of gender in society.

Gender is a conception that exists in the practice and processes of day-to-day life and social institutions. It has become a truth which distinguishes male from female or masculinity from femininity on the basis of well-defined and strictly followed assumptions. It is paradoxical that on one hand, gender is continuously changing and re-formulating in society, and on the other hand, it forms the basis and structure of social life. An individual's upbringing as well as surroundings determines the process of functioning according to gender. The formulation and maintenance of gender are evident in personal selves as well as in social communications. The reproduction of gender in society gives birth to the gendered structure of society. Gender is observable in our personalities, our cultures, our institutions in a complicated manner. In fact, gender leads to the construction of patterns of expectations for individuals, and is built in the vital components of society i.e. economy, family, and politics.

Gender can also be understood as a routine and methodical performance that keeps repeating itself, and relies on everyone performing gender. The competence of men and women performing gender depends upon the performance that men and women indulge in. Although the individual performs gender, yet gender is regarded as a distinct feature of social circumstances. Gender is both a result, and a principle for maintaining social order. The process of belonging to a particular gender is understood and standardized by the society, its values, in addition to religion, legal as well as scientific structure. It is not only about cultural aspects and traits that produce personal identities. Performing gender is associated with a particular sex and thus, can be regarded as a natural process. The distinctions between women

IJSS - Four International Journal of Social Sciences

Page 27

Vol. No. 04, Issue No. 16, December- 2017

A UGC Approved Journal No. 63712

ISSN - 2395-1133 (Print)

and men due to their sex transform them into fundamental and consistent characters. Since gender is instrumental in forming a structure of social life, gender statuses must be clearly distinguished. The resultant order of the society, thus, reflects fundamental and natural differences.

Incidentally, it can be contended that if we perform gender in a systematic manner, simultaneously sustain, construct and legitimise the institutional order that has a sex category as its basis. Thus, gender inequality is begotten by reproductive differences between women and men. Moreover, it cannot be said for sure that the subjugation of women as a result happened as an accident or as a deliberate action. One can however be sure of the fact that the gradual subordination of a particular gender has led to the formation of a group called women

4 | Page


Coordinator
NAAC
SMHS Government College
Sahibzada Ajit Singh Nagar


Harjeet Arjyal

Principal
SMHS Govt. College
Sahibzada Ajit Singh Nagar

International Journal of Advance Research in Science and Engineering
Volume No.07, Special Issue No.08, March 2018
www.ijarse.com

IJARSE
ISSN: 2319-8354

SUBCLASSES OF ANALYTIC FUNCTIONS RELATED TO SIGMOID FUNCTION

Gagandeep Singh¹, Gurcharanjit Singh², Harjinder Singh³

¹Department of Mathematics, Majha College for Women, Tarn-Taran (Punjab), India

²Department of Mathematics, Guru Nanak Dev University College, Chungh, Tarn-Taran (Punjab), India

³Assistant Director, Directorate of Public Instructions Punjab, Chandigarh (Punjab), India

ABSTRACT

In this paper, the authors investigate the initial coefficient bounds for some new subclasses of analytic functions related to Sigmoid function. Also the relevant connections to Fekete-Szegő inequality and Hankel determinant for these classes are briefly discussed. Our results serve as a new generalization in this direction.

Mathematics Subject Classification: 30C45, 33E99

Keywords: Analytic functions, Convex function, Sigmoid function, Starlike function, Subordination.

I INTRODUCTION AND PRELIMINARIES

The theory of special functions is significantly important to scientists and engineers. Though not with any specific definition but its applications extend to physics, computer etc. Recently, the theory of special functions has been overshadowed by other fields like real analysis, functional analysis, algebra, topology and differential equations.

There are various special functions but we shall concern with one of the activation function known as sigmoid function or simple logistic function. Activation function is an information process that is inspired by the biological nervous system such as brain processes information. It comprises of large number of highly interconnected processing element (neurons) working together to solve a specific task. The function works in similar way the brain does. it learns by examples and cannot be programmed to solve a specific task.

The sigmoid function of the form

$$h(z) = \frac{1}{1 + e^{-z}} \quad (1.1)$$

is differentiable and has the following properties:

- It outputs real numbers between 0 and 1.
- It maps a very large input domain to a small range of outputs.
- It never loses information because it is a one-to-one function.
- It increases monotonically.



Coordinator

NAAC

SMHS Government College
Sahibzada Ajit Singh Nagar



Principal

SMHS Govt. College

Sahibzada Ajit Singh Nagar



GLOBAL JOURNAL OF SCIENCE FRONTIER RESEARCH: I
MATHEMATICS AND DECISION SCIENCES
Volume 18 Issue 3 Version 1.0 Year 2018
Type: Double Blind Peer Reviewed International Research Journal
Publisher: Global Journals
Online ISSN: 2249-4626 & Print ISSN: 0975-5896

A New Subclass of Univalent Functions

By Gagandeep Singh, Gurcharanjit Singh & Harjinder Singh

Majha College for Women

Abstract- In this paper, a new subclass $\chi_1(A, B)$ of close-to-convex functions, defined by means of subordination is investigated. Some results such as coefficient estimates, inclusion relations, distortion theorems, radius of convexity and Fekete-Szegő problem for this class are derived. The results obtained here is extension of earlier known work.

Keywords: subordination, univalent functions, analytic functions, convex functions, close-to-convex, coefficient estimates, fekete-szegő problem.

GJSFR-F Classification: MSC 2010: 30C45



Strictly as per the compliance and regulations of:



© 2018. Gagandeep Singh, Gurcharanjit Singh & Harjinder Singh. This is a research/review paper, distributed under the terms of the Creative Commons Attribution-Noncommercial 3.0 Unported License <http://creativecommons.org/licenses/by-nc/3.0/>, permitting all non commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Coordinator

NAAC

SMHS Government College

Sahibzada Ajit Singh Nagar

Principal

SMHS Govt. College

Sahibzada Ajit Singh Nagar



Pardeep Rattan

Data mining: A library utility model

Authors Pardeep Rattan

Publication date 2019


Journal European Journal of Research

Volume 1

Pages 39-34

Description Documents, Users, Services, Finances, Human resources and Space are to library, what library is to an institution. Library perhaps is the only service agency that through its robust and helpful library system based on tools and techniques of information and communication technology is able to satisfy the information needs of a user across the globe 24x7x365 that comes out to be 61320 hours in an year. Libraries through Data mining (DM) techniques would be able to strengthen its managerial and decision support system where data is analysed from different perspectives which in turn would provide an edge to an organisation like library to serve their clients in a better way. An attempt has been made through this conceptual paper to identify the core library areas where data mining techniques can be applied to build a stronger serviceable library system for the maximum benefit of library users.

Total citations Cited by 5



Scholar articles Data mining: A library utility model
P Rattan - European Journal of Research, 2019
Cited by 5 Related articles

Coordinator
NAAC
SMHS Government College
Sahibzada Ajit Singh Nagar

Principal
SMHS Govt. College
Sahibzada Ajit Singh Nagar

THINK INDIA (Quarterly Journal)

ISSN: 0971-1180

Vol-22-Issue-4-October-December-2019

Economic Dilemma and Suicide Among Farmers In Punjab

Vipin Chaudhary¹ and (Dr.) Shruti Shourie²

¹Research Scholar, Department of Psychology, Panjab University Chandigarh.

(Email id- vipinchaudhary87@gmail.com)

²Associate Professor, D.A.V College, Sector- 10, Chandigarh.

Abstract

The current paper attempted to understand the economic dilemma and suicide among farmers in Punjab. For this purpose, a sample of 230 farmers was presented with nine questions, on whether they have taken any debt, reasons for high suicide rates, purpose of taking loans, reasons for not paying instalments of loan on time, and the biggest problem faced by farmers in Punjab in today's time. Farmers reported to close-ended questions, which had response options, such as, yes/no or other multiple responses. Data was analysed with regards to frequencies of responses for every question. These numerical proportions were further represented through a pie chart. Results are further discussed and suggestions are made based on the findings.

Keywords: Farmers, Punjab, Debt, Farmer Suicide, Indebtedness, Loans

Introduction

Agriculture is related to the word farming, and farming supports more than fifty per cent of the Indian population (Ashalatha & Rajeshwari, 2018). However, the contribution of agriculture in Gross Domestic Product (GDP) of India is declining steadily over the years due to myriad reasons related to social, political, and psychological aspects.

Despite Punjab's economic growth going forward, it boasts one of the highest unemployment rates in the country (Singh, 2018); therefore the farmers of Punjab, due to helplessness, rely upon farming as their only source of income. Farmers are not in a very rosy condition. They are facing distress due to factors such as cyclones, excessive or very little rainfall, drought, inflation, and floods. These factors have led to high mortality rates, revolts, protests, and frustration among farmers in the past few years (Ashalatha & Rajeshwari, 2018).

Farmers are accumulating high debt while lacking alternative sources of income.

THINK INDIA (Quarterly Journal)

ISSN 0971-1260
Vol-22-Issue-4-October-December-2019

**Relationship between Social Support and Resilience among Farmers
from Punjab**

Vipan Chaudhary¹ and (Dr.) Shruti Shourie²

¹Research Scholar, Department of Psychology, Punjab University Chandigarh.
(Email id- vipanchaudhary87@gmail.com)

² Associate Professor, D.A.V College, Sector- 10, Chandigarh.


Abstract

Farmers are socially isolated, working for long hours without any opportunity for social interaction, which can lead to loneliness and poor mental health (Skerratt, 2018). The present study aimed to assess the relationship between perceived social support and resilience among farmers. For this purpose, a sample of 230 farmers aged 35-50 years was selected from various rural areas of Punjab. Results revealed a significant positive relationship between perceived social support and resilience. This finding has implications for mental health counsellors in enhancing resilience among farmers by highlighting the role of social support through family counselling and awareness campaigns. Families of farmers need to be educated with regards to their role in providing social support to the farmers in the face of difficult times, and farmers need psychoeducation about seeking social support without hesitation or inhibitions.

Keywords: Farmers, Resilience, Social Support, Punjab


Introduction

Even though Punjab produces a maximum amount of crops and high-quality wheat than any other state, the state has faced 10,000 suicides in the past ten years (Khanna, 2019). In Punjab, 1.6 lakh farmers are facing severe stress as they are under debt (Bharti, 2018). 85.9 per cent of agricultural households in Punjab are living



Volume 2220, Issue 1

4 May 2020



3RD INTERNATIONAL
CONFERENCE ON
CONDENSED MATTER AND
APPLIED PHYSICS (ICC-
2019)

14–15 October 2019

Bikaner, India

< Previous Article

Next Article >

RESEARCH ARTICLE | MAY 04 2020

Investigating TiO₂ at two different calcination temperatures

Shilpy Bhullar, Shikha Gupta , Navdeep Goyal

Check for updates

+ Author & Article information

AIP Conference Proceedings 2220, 020065 (2020)

<https://doi.org/10.1063/5.0001662>

Share ▾

Tools ▾

Titanium dioxide was synthesized by sol-gel technique and calcined at two different temperatures 600°C (T600) and 800°C (T800). XRD and FTIR were performed on the two samples. XRD results showed that at 600°C, mixed-phase (anatase and rutile) was present with anatase peaks more prominent than the rutile. However, at 800°C rutile peaks became more prominent. The sizes of nanoparticles synthesized showed that the particle size increases as the temperature increases and as the phase changes from anatase to rutile. FTIR results showed that hydroxyl groups and surface adsorbed water were removed when the temperature was increased to 800°C. The characteristic stretching of Ti-O and Ti-O-Ti bonds, vibrations of -CH group and C=O group were retained by both T600 and T800.

Coordinator

NAAC

SMHS Government College
Sahibzada Ajit Singh Nagar

Principal

SMHS Govt. College
Sahibzada Ajit Singh Nagar

A Second Order Smoothing Penalty Function Algorithm for Constrained Optimization Problems

Turkish Online Journal of Qualitative Inquiry (TOJQI)

Volume 11, Issue 3, October 2020: 625-643

A Second Order Smoothing Penalty Function Algorithm for Constrained Optimization Problems

Darpan Sood¹, Dr. Amanpreet Singh², Dr. Rama³, Dr. Amrit Pal Singh⁴

¹Research Scholar, Department of Mathematics, Desh Bhagat University, Punjab, (India), 147301

²Assistant Professor, Department of Mathematics, GSSDGS Khalsa College, Patiala, Punjab, 147001

³Professor, Department of Mathematics, Desh Bhagat University, Punjab, (India), 147301

⁴Assistant Professor, Department of Mathematics, SMHS Government College, SAS Nagar, Punjab, 160055

Abstract The current paper introduces a second order smoothing technique for classical ϵ -exact penalty function in constrained optimization problems. Error calculations for optimum solution values for non-smoothed, smoothed penalty problem and for the original problem have been discussed in the paper. An algorithmic procedure for obtaining the solution is demonstrated and convergence is discussed.

Keywords Penalty Function, Smoothing, Error, Convergence, Constrained optimization problem

Introduction

The mathematical form of constraint optimization problem involves the introduction of certain terminology which should be known for better understanding of the topic. Let x be an n -dimensional vector given as $x = (x^1, x^2, \dots, x^n) \in R^n$. S be a subset of R^n . Let $f_0(x), f_1(x), \dots, f_m(x)$ are functions of x . The main problem in constrained optimization can be represented as

$$(1) \quad \text{Min } f_0(x)$$

$$\text{s.t. } f_j(x) \leq 0, j = 1, 2, 3, \dots, m$$

The function f_0 and $f_j, R^n \rightarrow R$ are continuous differentiable functions of second order. The function $f_0(x)$ is called the objective function. The vector function $f(x) = f_1(x), f_2(x), \dots, f_m(x)$ defined above is generally referred to as the functional constraints. The set S is called the basic feasible set. The set $Q = \{x \in S, f_j(x) \leq 0, j = 1, 2, 3, \dots, m\}$ is called the feasible set of the problem

(1). The set Q is assumed to be nonempty. The minimization problems can be classified as

1. Constrained Problems: $Q \subset R^n$

625

Taylor Francis Online

Home ▶ All Journals ▶ Asia Pacific Journal of Public Administration ▶ List of Issues ▶ Volume 43, Issue 4 ▶ Collaborative governance for urban sustainability

Asia Pacific Journal of Public Administration
Volume 43, 2021 - Issue 4


Submit an article Journal homepage

364 Views
2 CrossRef citations to date
5 Altmetric

Research Article
Collaborative governance for urban sustainability: implementing solar cities
Bharati Garg & Rohit Barach
Pages 236-257 | Published online: 28 May 2021

Download citation <https://doi.org/10.1080/23276655.2021.1925132>

Full Article Figures & data References Citations Metrics Reprints & Permissions Read this article



ABSTRACT

Today, when rapid urbanization has become an environmental concern all over the world, the concept of solar cities assumes a crucial role to play in realizing not only the sustainability potential of cities but also to transform urbanization into an opportunity. The Union Ministry of New and Renewable Energy (MNRE), Government of India in 2008 launched the Solar Cities Programme to promote alternate sources of energy and reduce dependency on conventional energy sources. Chandigarh was the first city to be declared as model solar city. This study attempts to empirically analyze the measures taken to make Chandigarh a model solar city while focusing on adoption of roof-top Solar Photovoltaic (SPV). The data collected was analysed and interpreted using percentages. Descriptive analysis of the responses of government officials, residents, NGOs and private empanelled agencies was done. Findings revealed that the whole idea of participatory local planning was sidelined, awareness generation remained piecemeal, no formal co-ordination mechanism was established and yet empirical targets were not only achieved but surpassed. However, the implementation failed to convince residents to shift towards solar energy, jeopardizing the long-term sustainability of the solar city plan.

KEYWORDS: Sustainability, Sustainable Urban Development, Sustainable Cities, Solar City, Public Policy, Collaborative Governance


Coordinator
NAAC
SMHS Government College
Sahibzada Ajit Singh Nagar


Principal
SMHS Govt. College
Sahibzada Ajit Singh Nagar

RSC Advances

ROYAL SOCIETY OF CHEMISTRY

PAPER

View Article Online
DOI: 10.1039/C1AY01000A

Check for updates

Cite this: RSC Adv., 2021, 11, 30343

Rapid green-synthesis of TiO₂ nanoparticles for therapeutic applications

Shilpy Bhullar^a, Navdeep Goyal^a and Shikha Gupta^{a,b*}

Nanoparticles (NPs) with sizes ranging from 2 nm to 1 μm find various applications in the field of therapeutics. Moreover, if eco-friendly methods are opted for the synthesis of low-cost, stable and less toxic NPs, then that's a huge success. Titanium dioxide nanoparticles (TiO₂ NPs) have been extensively studied for their use in medical implants, photodynamic therapy, drug delivery, biosensing and as antimicrobial agents. The present study reports the green-synthesis of TiO₂ NPs for the first time using extracts of black pepper (*Piper nigrum*), eonlander (*Canarium sativum*) and clove (*Syzygium aromaticum*). All three samples of TiO₂ NPs were synthesized via a modified sol-gel method under similar environmental conditions. Similar treatments were given to the samples. The procedure adopted for the synthesis ensures the use of non-toxic materials, no production of toxic by-products and rapid synthesis of the TiO₂ NPs. The NPs were characterized by X-ray diffraction, high resolution transmission electron microscopy, energy dispersive spectroscopy, field emission scanning electron microscopy and selected area electron diffraction which confirmed the formation, morphology, crystallinity and size of the TiO₂ NPs. These characterizations displayed the similarity index of all three samples. However, photoluminescence and vibrating sample magnetometer studies highlighted the differences among the three samples. All three samples of NPs obtained had a size range of 5–20 nm. Further, the findings showed that different plant extracts result in TiO₂ NPs with moderately different characteristics. Furthermore, the samples were analysed for their drug-encapsulation efficiency using UV-visible spectrophotometry. Among all three samples, the NPs synthesised using black pepper exhibited the maximum encapsulation efficiency. The study concludes that the plant's bio-profile is responsible for bringing about changes in the traits of the resulting nanoparticles. Thus, the extracts from different plants have the ability to manipulate the properties of the synthesized NPs. These findings can help to understand the role and importance of the plants in synthesizing NPs for biomedical applications. A further detailed study in this field can help researchers to understand the influence of the plant's biochemistry in shaping the NPs.

Received 23rd July 2021
Accepted 27th August 2021
DOI: 10.1039/C1AY01000A
rsc/advances

1 Introduction

Today the world is moving towards adopting eco-friendly measures for sustainable development. Nanoparticles (NPs) are now slowly capturing the global market because of their versatility. But as everything comes with a flaw, these NPs also result in biohazards. During their synthesis, chemicals are used which often lead to many toxic by-products. As a remedy to this bizarre scenario, the green synthesis approach is fascinating to researchers these days. The green synthesis technique involves the maximum use of biomass to retrieve the NPs. However, the term 'green' is not restricted to plants alone. Various fungi-mediated and bacteria-mediated synthesis also come under

the green synthesis procedure. Thus, biomass, in general, is being extensively utilised for environment-friendly synthesis techniques. When using plants, the extracts of their roots, leaves, stems, seeds, flowers or fruits, can be used. Organic polymers are the building blocks of the plants and these include starch, chitin, cellulose, hemicellulose, lignin and various resins. Whenever plant extracts are used, the organic polymers and the biomolecules present in them, are responsible for their characteristic behaviour. Cellulose is the most abundant organic polymer and is a polysaccharide present in the primary cell wall of plants and many forms of algae. Lignocellulose biopolymers nourish the cell wall of the plants and they consist of cellulose, hemicellulose and lignin. Lignin constitutes around 10 percent of the lignocellulose biomass and contains a large number of phenylpropanoids. It has the potential to replace petroleum and its depolymerization offers remarkable possibilities for producing high-quality chemicals.^{1,2} This is a step towards modern day ecofriendly synthesis procedures.

^aDepartment of Physics, Centre of Advanced Study in Physics, Punjab University, Chandigarh-160014, India

^bDepartment of Physics, Government Gaurish Datta Sanskrit Dharma College, Sector-32C, Chandigarh-160033, India. E-mail: shikha.gupta@gsdcl.ac.in



Coordinator
NAAC
SMHS Government College
Sahibzada Ajit Singh Nagar

Harjeet Goyal

Principal
SMHS Govt. College
Sahibzada Ajit Singh Nagar

FEKETE-SZEGÖ INEQUALITY FOR CERTAIN CLASSES OF CLOSE-TO-CONVEX FUNCTIONS

GAGANDEEP SINGH, GURCHARANJIT SINGH, HARINDER SINGH

ABSTRACT. Close-to-convex functions and quasi-convex functions are of great importance in geometric function theory. In the present investigation, the authors study the subclasses C_1 of close-to-convex functions and the subclasses C^* and C_1^* of quasi-convex functions in the open unit disc $E = \{z : |z| < 1\}$. The sharp upper bounds of the functional $|\alpha_2 - \mu\alpha_1^2|$, μ real, for the functions of the form $f(z) = z + \sum_{n=2}^{\infty} a_n z^n$ belonging to these classes are provided. This work will pave the way to investigate the upper bound of the Fekete-Szegő functional for some other subclasses of close-to-convex and quasi-convex functions.

1. INTRODUCTION

Let A denote the class of functions of the form

$$f(z) = z + \sum_{n=2}^{\infty} a_n z^n \quad (1)$$

which are analytic in the unit disc $E = \{z : |z| < 1\}$. Let S be the class of functions of the form (1) which are analytic univalent in E .

We shall concentrate on the coefficient problem for the class S and certain of its subclasses. In 1916, Bieberbach [3] proved that $|a_2| \leq 2$ for $f(z) \in S$ as a corollary to an elementary area theorem. He conjectured that, for each function $f(z) \in S$, $|a_n| \leq n$; equality holds for the Koebe function $k(z) = z/(1-z)^2$, which maps the unit disc E onto the entire complex plane minus the slit along the negative real axis from $-\frac{1}{2}$ to $-\infty$. De Branges [5] solved the Bieberbach conjecture in 1984. The contribution of Löwner [10] in proving that $|a_3| \leq 3$ for the class S was huge.

With the known estimates $|a_2| \leq 2$ and $|a_3| \leq 3$, it was natural to seek some relation between a_2 and a_3^2 for the class S . This thought prompted Fekete and Szegő [6] and they used Löwner's method to prove the following well-known result for the class S :

1991 Mathematics Subject Classification. 30C45, 30C50.

Key words and phrases. Univalent functions, starlike functions, convex functions, close to convex functions, bounded functions.

Submitted April 30, 2020. Revised May 24, 2020.



Coordinator

NAAC

SMHS Government College

Sahibzada Ajit Singh Nagar



Principal

SMHS Govt. College

Sahibzada Ajit Singh Nagar



Australian Government
IP Australia

CERTIFICATE OF GRANT INNOVATION PATENT

Patent number: 2021107060

The Commissioner of Patents has granted the above patent on 17 November 2021, and certifies that the below particulars have been registered in the Register of Patents.

Name and address of patentee(s):

Amanpreet Singh of GSDS Khalsa College Patiala Punjab India
Darpan Sood of SGTB Khalsa College Anandpur Sahib Punjab India
Amrit Pal Singh of SMHS Government College SAS Nagar Punjab India

Title of invention:

A SMOOTHING TECHNIQUE FOR SQUARE ROOT EXACT PENALTY FUNCTION IN CONSTRAINED OPTIMIZATION

Name of inventor(s):

Singh, Amanpreet; Sood, Darpan and Singh, Amrit Pal

Term of Patent:

Eight years from 24 August 2021

NOTE: This Innovation Patent cannot be enforced unless and until it has been examined by the Commissioner of Patents and a Certificate of Examination has been issued. See sections 120(1A) and 129A of the Patents Act 1990, set out on the reverse of this document.

Priority details:

Number	Date	Filed with
202111035302	5 August 2021	IN



Coordinator
NAAC
SMHS Government College
Sahibzada Ajit Singh Nagar


Principal
SMHS Govt. College
Sahibzada Ajit Singh Nagar



Volume 2352, Issue 1

5 August 2021



ADVANCED MATERIALS
AND RADIATION PHYSICS
(AMRP-2020): 5th National
e-Conference on Advanced
Materials and Radiation
Physics

9–11 November 2020

Longowal, India

< Previous Article

Next Article >

RESEARCH ARTICLE | AUGUST 05 2021

Factors Influencing synthesis of titanla nanoparticles – A short review

Shilpy Bhullar, Navdeep Goyal, Shikha Gupta

Check for updates

+ Author & Article Information

AIP Conference Proceedings 2352, 040036 (2021)

<https://doi.org/10.1063/5.0052996>

Share

Tools

There is a plethora of factors which affect the synthesis of Titanium Dioxide (TiO_2) nanoparticles (NPs). There is a great deal of benefits if we know how to optimize different factors to tune the properties of NPs. Out of many factors like precursor, pH, mode of synthesis, time, pressure, etc. temperature and reagents play a great role in modulating NPs. Even though, many researchers are making nanoparticles by following different approaches, not much is known about how and why do these factors actually affect the NPs during synthesis. The present review highlights the works where temperature and nature of reagents or solvents pose a significant contribution in the manufacturing of stable nanoparticles. Moreover, possible explanations would be put forth describing the observed behaviour.

Coordinator

NAAC

SMHS Government College

Sahibzada Ajit Singh Nagar

Principal

SMHS Govt. College


Sahibzada Ajit Singh Nagar



AIP Conference Proceedings

HOME BROWSE FOR AUTHORS ▾ FOR ORGANIZERS ▾ ABOUT ▾

Volume 2357, Issue 1
9 May 2022



NATIONAL CONFERENCE
ON ADVANCES IN APPLIED
SCIENCES AND
MATHEMATICS: NCASM-20
24-25 September 2020
Rajpura, India

[← Previous Article](#) [Next Article →](#)

[tcp/issue/2357/1](#)

RESEARCH ARTICLE | MAY 09 2022

Titanium dioxide nanoparticles synthesized using different reagents

Shilpy Bhullar, Navdeep Goyal, Shikha Gupta

[Check for updates](#)

+ Author & Article Information

AIP Conference Proceedings 2357, 050015 (2022)

<https://doi.org/10.1063/5.0080593>


[Share](#)

[Tools](#)

Titanium dioxide nanoparticles (TiO₂ NPs) have been synthesized by researchers all over the world by using different methods and different reagents. Sol-gel technique is, by far, the most suitable and most convenient method to synthesize nanoparticles. In the present study, TiO₂ nanoparticles were manufactured following the sol-gel route using Glacial Acetic Acid (GTDNPs) as the main reagent. In our previous study, we had synthesized titanium dioxide nanoparticles using Ethanol (ETDNPs) as one of the major constituents. An attempt has been made to compare the presently manufactured GTDNPs and the previously synthesized ETDNPs. Moreover, similar treatment was given to both the samples. The findings indicated that similar XRD patterns were obtained in both cases which confirmed the formation of TiO₂ NPs. Although, rutile peaks were more dominant than the anatase peaks in GTDNPs, the opposite was observed in ETDNPs and the crystallite size obtained by Scherrer's equation was found to be greater in ETDNPs than in GTDNPs. This is a very crucial result to understand the role of different reagents played on the size of nanoparticles so formed.

Coordinator
NAAC
SMHS Government College
Sahibzada Ajit Singh Nagar

Principal
SMHS Govt. College
Sahibzada Ajit Singh Nagar



Coordinator
NAAC
SMHS Government College
Sahibzada Ajit Singh Nagar

Harjeet Goyal

Principal
SMHS Govt. College
Sahibzada Ajit Singh Nagar

Journal of Rajasthan Academy of Physical Sciences

ISSN: 0972-6306; URL: <http://raops.org.in>

Vol.21, No.1&2, January-June, 2022, 65-70

COEFFICIENT INEQUALITY FOR A COMBINED SUBCLASS OF VARIOUS CLASSES OF REGULAR FUNCTIONS

Gurmeet Singh

Department of Mathematics, Khalsa College, Patiala

E-mail: meetgur111@gmail.com

Harjinder Singh (Corresponding Author) SMHS Govt. College, Mohali

E-mail: harjindpreet@gmail.com

Misha Rani

Research Fellow, Punjabi University, Patiala

E-mail: mishagargsamana@gmail.com

Abstract. Here, we take functions of the type $f(z) = z + \sum_{k=2}^{\infty} a_k z^k$ and solve the Fekete – Szegő inequality for a new class of analytic functions.

2010 Mathematics Subject Classification: 30C45, 30C50.

Keywords: Fekete – Szegő Inequality, Starlike functions, Bounded analytic functions and concept of subordination.

1. Introduction

In this paper we define an inequality called Fekete – Szegő Inequality for a new class of analytic functions. This is an inequality which relates to those coefficients which are related to univalent analytic functions [8],[16]. M. Fekete and G. Szego proved this inequality in 1933[5]. It originates from bieberbach conjecture([6], [13], [14], [15]), which was given by Bieberbach [2] in 1916 but finally proved by him [3] in 1985.

Firstly, let us discuss some classes and some basic results :

Let A be the family of functions f of the form $f(z) = z + \sum_{k=2}^{\infty} a_k z^k$, having conditions of normalisation $f(0) = 0, f'(0) = 1$; analytic in open unit disc $E = \{z \in C: |z| < 1\}$.

Let S be the family of functions f univalent in the open disk $\{z \in C: |z| < 1\}$ with conditions

$$f(0) = 0, f'(0) = 1; f(z) = z + \sum_{k=2}^{\infty} a_k z^k.$$

Any function f belonging to the class A is said to be a Starlike function if $f(E)$ is starlike domain with respect to the origin and this class is denoted by S^* [1]. The essential condition for this class as given by Duren [4], is $Re \left(\frac{zf'(z)}{f(z)} \right) > 0; z \in E$, and $S^*(\phi)$ be the class of functions in $f \in S$, for which $\frac{zf'(z)}{f(z)} < \phi(z)$, given by Ma and Minda [10].



Coordinator

NAAC

SMHS Government College
Sahibzada Ajit Singh Nagar



Principal

SMHS Govt. College
Sahibzada Ajit Singh Nagar